arXiv:1208.3124v1 [cs.CG] 14 Aug 2012

ON THE COMPUTATION OF ZONE AND DOUBLE ZONE DIAGRAMS

DANIEL REEM

Abstract. A zone diagram is a relatively new concept related to computational geometry and fixed point theory. More precisely, it is a variation of the notion of the Voronoi diagram and formally it is a fixed point of a certain mapping. Neither its uniqueness nor its existence are obvious in advance. It has been studied by several authors, starting with T. Asano, J. Matoušek and T. Tokuyama who considered the Euclidean plane with point sites. Computation of zone diagrams is a challenging task and in a continuous setting it has been addressed (briefly) only by these authors in the above setting. We consider the question of computation of zone diagrams in finite dimensional strictly convex spaces with general positively separated closed sites. We show that a generalization of the iterative algorithm suggested by these authors converges to a double zone diagram, a concept closely related to a zone diagram whose existence is known in general. In many cases a zone diagram can be obtained from the resulting double zone diagram. We are able to compute approximately the corresponding iterations and the resulting (double) zone diagram using a new algorithm which enables the computation of Voronoi diagrams in a general setting. Unexplained interesting phenomena are discussed too.

1. Introduction

1.1. **Background.** A zone diagram is a relatively new concept related to geometry and fixed point theory. In order to understand it better, consider first the more familiar concept of the Voronoi diagram. Given a set X, a distance function d, and a collection of subsets $(P_k)_{k\in K}$ in X (called the sites or the generators), we associate with each site P_k a corresponding Voronoi cell, that is, the set R_k of all $x\in X$ whose distance to P_k is not greater than its distance to the union of the other sites P_j, $j\neq k$. On the other hand, in the case of a zone diagram $R=(R_k)_{k\in K}$ we associate with each site P_k the set R_k of all $x\in X$ whose distance to P_k is not greater than its distance to the union of the other sets R_j, $j\neq k$. Figures 1 and 2 show the Voronoi and zone diagrams, respectively, corresponding to the same ten singleton sites in the Euclidean plane.

At first sight it seems that the definition of a zone diagram is circular, because the definition of each R_k depends on R_k itself via the definition of the other cells R_j, $j\neq k$. On second thought, we see that, in fact, a zone diagram is defined to be a fixed point of a certain mapping (called the Dom mapping), that is, a solution of the fixed point equation $R=\mathrm{Dom}(R)$. While the Voronoi diagram is explicitly defined and, hence, its existence and uniqueness are obvious, neither the existence nor the uniqueness of a zone diagram are obvious in advance. In addition, even if some existence (or uniqueness)

2010 *Mathematics Subject Classification.* 47H10, 68U05, 46B20.

Key words and phrases. Algorithm, computation, double zone diagram, fixed point, iteration, strictly convex normed space, Voronoi diagram, zone diagram.



FIGURE 1. The Voronoi diagram of 10 point sites in a square in the Euclidean plane.



FIGURE 2. The zone diagram of the same 10 points as in Figure 1.

results are proved, one still faces the problem of finding algorithms for computing zone diagrams.

The concept of a zone diagram was first defined and studied by T. Asano, J. Matoušek and T. Tokuyama in [2, 3] in the case where (X, d) was the Euclidean plane, each site P_k was a single point, and all these (finitely many) points were different. They proved the existence and uniqueness of a zone diagram in this case, and also suggested a natural iterative algorithm for approximating it. Their proofs and methods rely heavily on the above setting.

Later, more general existence results were obtained. For instance, in [22] it was shown that a zone diagram of two general sites in any metric space always exists. In fact, the proof holds in a more general setting called m-spaces, in which X is an arbitrary nonempty set and the "distance" function d should only satisfy the condition $d(x, x) \leq d(x, y)\ \ \forall x, y \in X$ and can take any value in the interval $[-\infty, \infty]$. Simple examples given there show that in general uniqueness of the zone diagram does not necessarily hold. In [15] it was shown, in particular, that a zone diagram of any finite number of sites exists, assuming these sites are compact and positively separated (that is, there is a positive lower bound on the distance between any pair of them) and that they are located in a large compact subset of $\mathbb{R}^n$ with a strictly convex norm (the exact result is more general). Another existence (and also uniqueness) result was discussed in [12]. Here the setting was the n-dimensional Euclidean space $\mathbb{R}^n$, or, more generally, finite dimensional normed spaces which are both strictly convex and smooth [12]. The sites were again positively separated.

The existence of a zone diagram is not known in general. However, as proved in [22], the existence of a closely related concept called a double zone diagram is indeed known in general (in any m-space, for infinitely many arbitrary sites). Double zone diagrams were introduced in [22] as a means for proving the existence of a zone diagram, partly because any zone diagram is a double zone diagram. Their existence was shown in a nonconstructive way (using the Knaster-Tarski fixed point theorem [14, 25]), and, as a matter of fact, it was shown that there exist a least and a greatest double zone diagrams. The importance of double zone diagrams to the computation of zone diagrams will become clear later. Zone diagrams are also related to other concepts, such as the concepts of trisectors [1, 2, 4, 7], k-sectors and k-gradations [10], and territory diagrams [8].

One of the main challenges regarding zone diagrams is their computation. In a continuous setting this task has been addressed so far only by Asano, Matoušek, and Tokuyama [3] in the case of the Euclidean plane with finite number of distinct points. The discussion about an actual method for approximate computing was very brief with almost no theoretical or practical details. In spite of the fact that several pictures were presented in [3], the only clue regarding the method of computation was mentioned briefly at the bottom of page 1184. It was written there that one can use convex polygons with many sides for approximating the components of these sequences, and that each iteration is computationally demanding. This (approximate) computation was restricted to the above setting. As will be explained in Section 3, one of the main reasons for this restriction was the lack of a method for computing Voronoi diagrams in a general setting. Another reason is the lack of any known representation of the boundaries of the involved regions (conjectured to be non-algebraic in many cases).

In a discrete setting (X is a finite set of points) there has been a limited discussion in the case of two sites: a point and a "line segment" or a "curve" in the digital Euclidean plane [6] (it was claimed there on the last page that the algorithm can be generalized to finitely many line segments but no details were given) and two finite sites in any finite m-space [22]. The actual computation in the first case is restricted to the above setting and no convergence analysis was given, and there were only clues regarding the time complexity. In the second case the actual computation is by brute force. In addition, it is not clear in which sense the resulting discrete zone diagrams approximate the continuous ones, although intuitively they may approximate them with respect to the Hausdorff distance. The computation of double zone diagrams was discussed briefly in [22] only in a discrete setting (finitely many finite sites in a finite m-space).

1.2. **Contribution of this paper.** We consider the question of computing zone diagrams in finite dimensional strictly convex spaces with general positively separated closed sites. We show that a generalization of the iterative algorithm suggested by Asano, Matoušek, and Tokuyama converges to the least and the greatest double zone diagrams. In various cases a zone diagram can be obtained from the resulting double zone diagrams, and as a matter of fact, in many of these cases (such as the Euclidean case) the limit is the unique zone diagram. We are able to compute approximately the corresponding iterations and the resulting (double) zone diagram using a new algorithm for computing Voronoi diagrams [19, 20] which enables their approximate computation in any normed space (possibly infinite dimensional) and with sites of a general form. Many pictures of (double) zone diagrams or their approximating sets produced by the method described here are given. It is interesting to note that the main ingredient in the computation of each iteration, namely the algorithm for computing Voronoi diagrams, enables the computation of each cell independently of the other ones and hence it and the above iterative algorithm can be easily parallelized. The Voronoi algorithm [19] can also help regarding the computation of k-sectors [10] and territory diagrams

[8], but these issues will not be dealt with here (see [10, p. 719] for a short related discussion).

1.3. **The structure of this paper.** In Section 2 we present the basic definitions and notation. In Section 3 we discuss qualitatively the iterative algorithm. In Section 4 we present the main convergence result. In Section 5 we explain how the corresponding iterations are actually computed and we also we briefly discuss the time complexity of the algorithm, as well as other topics. The proof of the main result is given in Section 6. We conclude the paper with Section 7, which contains several interesting questions and unexplained phenomena.

2. NOTATION AND DEFINITIONS

In this section we present our notation and basic definitions. We consider a nonempty convex subset X in some normed space $(\widetilde{X}, |\cdot|)$. We assume that X is not a singleton, for otherwise everything is trivial. We note that the definitions given below can be generalized almost word by word to the case where (X, d) is a general metric space and sometimes even beyond (m-spaces). We denote by $\overline{A}$, $\text{Int}(A)$ and ∂A the closure, interior, and the boundary, respectively, of the set A .

Throughout the text we will make use of tuples, the components of which are sets (which are subsets of the given set X). Every operation or relation between such tuples, or on a single tuple, is done component-wise. Hence, for example, if $K \neq \emptyset$ is a set of indices, and if $R = (R_k)_{k\in K}$ and $S = (S_k)_{k\in K}$ are two tuples of sets, then $R \bigcap S = (R_k \bigcap S_k)_{k\in K}$, $\overline{R} = (\overline{R_k})_{k\in K}$, and $R \subseteq S$ means $R_k \subseteq S_k$ for each $k \in K$. When R is a tuple, the notation $(R)_k$ is the k-th component of R, i.e, $(R)_k = R_k$.

Definition 2.1. *Given two nonempty sets $P, A \subseteq X$, the dominance region* $\text{dom}(P, A)$ *of P with respect to A is the set of all $x \in X$ the distance of which to P is not greater than its distance to A, that is,*

$$\text{dom}(P, A) = \{x \in X : d(x, P) \leq d(x, A)\}.$$

Here $d(x, A) = \inf\{d(x, a) : a \in A\}$.

Definition 2.2. *Let K be a set of at least 2 elements (indices), possibly infinite. Given a tuple $(P_k)_{k\in K}$ of nonempty subsets $P_k \subseteq X$, called the generators or the sites, the Voronoi diagram induced by this tuple is the tuple $(R_k)_{k\in K}$ of nonempty subsets $R_k \subseteq X$, such that for all $k \in K$,*

$$R_k = \text{dom}(P_k, \bigcup_{j\neq k} P_j) = \{x \in X : d(x, P_k) \leq d(x, P_j)\ \forall j \neq k,\ j \in K\}.$$

In other words, each R_k, called a Voronoi cell or a Voronoi region, is the set of all $x \in X$ whose distance to the site P_k is not greater than its distance to any other site P_j, $j \neq k$.

Definition 2.3. *Let K be a set of at least 2 elements (indices), possibly infinite. Given a tuple $(P_k)_{k\in K}$ of nonempty subsets $P_k \subseteq X$, a zone diagram with respect to that tuple is a tuple $R = (R_k)_{k\in K}$ of nonempty subsets $R_k \subseteq X$ such that*

$$R_k = \text{dom}(P_k, \bigcup_{j\neq k} R_j) \quad \forall k \in K.$$

In other words, if we define $X_k = \{C : P_k \subseteq C \subseteq X\}$*, then a zone diagram is a fixed point of the mapping* $\mathrm{Dom} : \prod\limits_{k\in K} X_k \to \prod\limits_{k\in K} X_k$*, defined by*

$$\mathrm{Dom}(R) = (\mathrm{dom}(P_k, \bigcup_{j\neq k} R_j))_{k\in K}. \tag{1}$$

a tuple $R = (R_k)_{k\in K}$ *is called a double zone diagram if it is a fixed point of the second iteration* $\mathrm{Dom} \circ \mathrm{Dom}$*, i.e.,* $R = \mathrm{Dom}^2(R)$*.*

Note that if R is a zone diagram, then it is a double zone diagram as implied by iterating the equation $R = \mathrm{Dom}(R)$ with Dom. We now recall the definition of strictly convex spaces.

Definition 2.4. *A normed space* $(\widetilde{X}, |\cdot|)$ *is said to be strictly convex if* $|x+y| < |x| + |y|$ *whenever* x *and* y *are any elements in* $\widetilde{X}$ *which are not on the same ray, i.e.,* $x \neq 0, y \neq 0$*, and* $x/|x| \neq y/|y|$*.*

A well known equivalent condition for strict convexity is that the unit sphere of the space does not contain line segments. Equivalently, for all $x, y \in \widetilde{X}$ satisfying $|x| = |y| = 1$ and $x \neq y$, the inequality $|(x+y)/2| < 1$ holds. The d-dimensional Euclidean space $\mathbb{R}^d$, or more generally, inner product spaces, and the sequence spaces ℓ_p, $p \in (1, \infty)$, are all examples of strictly convex spaces. On the other hand, $\mathbb{R}^d$ with the ℓ_1 or ℓ_∞ norms are typical examples of spaces which are not strictly convex. See [16, 18] for more information regarding strictly convex spaces.

We finish this section by recalling the definition of the Hausdorff distance.

Definition 2.5. *Given two nonempty sets* $A_1, A_2 \subseteq X$*, the Hausdorff distance between them is defined by*

$$D(A_1, A_2) = \max\{\sup_{a_1\in A_1} d(a_1, A_2), \sup_{a_2\in A_2} d(a_2, A_1)\}.$$

Recall that the Hausdorff distance is different from the usual distance between two sets which is defined by $d(A_1, A_2) = \inf\{d(a_1, a_2) : a_1 \in A_1, a_2 \in A_2\}$.

3. A QUALITATIVE DESCRIPTION OF THE ALGORITHM

As mentioned before, a tuple $R = (R_k)_{k\in K}$ of cells (subsets) is a zone diagram if it satisfies the fixed point equation $R = \mathrm{Dom}(R)$. A common and natural approach in fixed point theory for the computation of a fixed point of a given mapping f is by iterations [9, 13]. One starts with some point y_0 in the space Y on which f is defined, and starts iterating f. A sequence $y_1 = f(y_0), \ldots, y_{n+1} = f(y_n), \ldots$ is generated, and one hopes that it converges in some sense to a fixed point of f.

Returning to our setting, the given mapping is the Dom mapping. The given space on which it is defined is $Y = \prod_{k\in K} X_k$, where $X_k = \{C : P_k \subseteq C \subseteq X\}$. In other words, Y is the collection of all tuples (vectors) whose k-th component is a subset C of the given world X such that C contains the site P_k. As a result of the above, a natural choice for the starting point $y_0 \in Y$ is the collection $(P_k)_{k\in K}$ of the given sites.

It turns out that it is convenient to denote $I^{(0)} := (P_k)_{k\in K} = y_0$, $O^{(0)} := \mathrm{Dom}(I^{(0)})$, and define inductively the inner sequence $I^{(n)} := \mathrm{Dom}(O^{(n-1)})$ and the outer one $O^{(n)} := \mathrm{Dom}(I^{(n)})$ for any integer $n \geq 1$. The reason behind this choice is explained



FIGURE 3. The boundaries of $I^{(2)}$ and $O^{(2)}$ in a square in $(\mathbb{R}^2,\ell_2)$. The sites are as in Figure 2.



FIGURE 4. The setting of Figure 3, but with $I^{(3)}$ and $O^{(3)}$.

in the following simple lemma, which was essentially observed in [3] and a little bit in [20, 22].

Lemma 3.1. *Let $(P_k)_{k\in K}$ be a tuple of nonempty and closed subsets in X.*

(a) Dom *is antimonotone, i.e.,* $\mathrm{Dom}(R)\subseteq \mathrm{Dom}(S)$ *whenever* $S\subseteq R$*;* Dom^2 *is monotone, that is,* $R\subseteq S\Rightarrow \mathrm{Dom}^2(R)\subseteq \mathrm{Dom}^2(S)$.

(b) The sequence $(I^{(n)})_{n=0}^{\infty}$ is increasing, $(O^{(n)})_{n=0}^{\infty}$ is decreasing and we have

$$I^{(n)}\subseteq \bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P_k)_{k\in K}\subseteq \bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}\subseteq O^{(n)} \tag{2}$$

for each nonnegative n. In addition, if R is a zone or double zone diagram in X, then

$$I^{(n)}\subseteq \bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P_k)_{k\in K}\subseteq R\subseteq \bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}\subseteq O^{(n)}\quad \forall n\geq 0. \tag{3}$$

Figures 3 and 4 show $I^{(2)},O^{(2)}$ and $I^{(3)},O^{(3)}$ respectively, in a square in $(\mathbb{R}^2,\ell_2)$. The sites are as in Figure 2. For another example, see Figure 11.

From the lemma we see that the goal of $I^{(n)}$ is to approximate the zone diagram from below, and the goal of $O^{(n)}$ is to approximate it from above. However, at this stage several difficulties arise. First, it is not clear that these sequences converge, and if they do converge, it is not clear that both of them converge to the same limit and that the limit of at least one of them is indeed a zone diagram.

Second, even if a convergence result is obtained, one faces the problem of the actual computation of $I^{(n)}$ and $O^{(n)}$ for $n\geq 1$. Even for $n=0$ this is not a simple task since $O^{(0)}=\mathrm{Dom}(I^{(0)})=(\mathrm{dom}(P_k,\bigcup_{j\neq k}P_j))_{k\in K}$ is the Voronoi diagram of the sites $(P_k)_{k\in K}$, and hence, for sites of a general form, or sites which possibly form a degenerated configuration, or for space with a general norm, one has to find algorithms which enable the computation of Voronoi diagrams in such a setting, and most of the familiar algorithms for computing Voronoi diagrams (see e.g., [5, 17, 24] for some reviews) are not able to achieve this task (or, in some cases, they are too complicated or too slow). Even if one restricts oneself to the familiar case of the Euclidean plane with point sites, one still faces problems starting from the iteration $n = 1$. This is because one has to know how to compute the components of $\mathrm{Dom}(S)$ for a given tuple $S = (S_k)_{k\in K}$, and hence one has to know a method for computing $\mathrm{dom}(P_k, \bigcup_{j\neq k} S_j)$, i.e., the Voronoi cell of P_k with respect to the set $\bigcup_{j\neq k} S_j$. Unfortunately, when $S = I^{(n)}$, $n \geq 1$, or $S = O^{(n)}$, $n \geq 0$, the components of S are general sets, and again, it seems that most of the familiar algorithms for computing Voronoi diagrams are not helpful here.

The sequences $(I^{(n)})_{n=0}^{\infty}$ and $(O^{(n)})_{n=0}^{\infty}$ were introduced in [3] in the case of the Euclidean plane with finitely many distinct point sites. It was shown that $\overline{\bigcup_{n=0}^{\infty} I^{(n)}} = \bigcap_{n=0}^{\infty} O^{(n)}$ and that this tuple is the unique zone diagram. Although this is not a pure convergence result (no limits) and although no error estimates for the level of approximation were given, one still has the plausible result that $(I^{(n)})_{n=0}^{\infty}$ increases to a zone diagram and $(O^{(n)})_{n=0}^{\infty}$ decreases to it. As for the actual computation of $I^{(n)}$ and $O^{(n)}$, it was already mentioned in the introduction that very few details were given there.

4. The convergence theorem

As mentioned in Section 3, it was proved in [3] that in the case of the Euclidean plane with point sites one has the equality $m := \overline{\bigcup_{n=0}^{\infty} I^{(n)}} = \bigcap_{n=0}^{\infty} O^{(n)} := M$, and $m = M$ is the unique zone diagram. As Theorem 4.1 below shows, in the more general setting of strictly convex spaces with possibly infinitely many general positively separated sites one has the weaker result that $m = \mathrm{Dom}(M)$ and $M = \mathrm{Dom}(m)$, and both m and M are double zone diagrams rather than zone diagrams (and they are not necessarily equal). In addition, Theorem 4.1 also discusses another way to obtain a zone diagram in the special case of two sites.

A corollary to the theorem ensures that when the world X is compact, then $(I^{(n)})_{n=0}^{\infty}$ and $(O^{(n)})_{n=0}^{\infty}$ converge to m and M respectively with respect to the Hausdorff distance. As a result, since in practice X is taken to be compact, if $I^{(n)}$ and $O^{(n)}$ are shown experimentally to be almost the same for some n, then one has a good approximation to both m and M. Because in this case it is known that a zone diagram exists [15], one has a good approximation to this zone diagram, and in fact this also shows that probably this zone diagram is unique and coincides with both m and M. In all of the pictures presented in this paper this is what we obtained. Another corollary to the above ensures that whenever the least and the greatest double zone diagrams coincide, then $m = M$ and they both coincide with the unique zone diagram. This is true in particular in the Euclidean case.

Theorem 4.1. *Let X be a convex set in a finite dimensional strictly convex space and let $P = (P_k)_{k\in K}$ be a tuple of nonempty closed sets in X with the property that*

$$\inf\{d(P_k, P_j) : j, k \in K,\, j \neq k\} > 0.$$

Let

$$M = \bigcap_{\gamma=0}^{\infty} \mathrm{Dom}^{2\gamma+1}(P) = \bigcap_{n=0}^{\infty} O^{(n)}, \quad m = \overline{\bigcup_{\gamma=0}^{\infty} \mathrm{Dom}^{2\gamma}(P)} = \overline{\bigcup_{n=0}^{\infty} I^{(n)}}.$$

Then

$$m = \mathrm{Dom}(M) \quad \text{and} \quad M = \mathrm{Dom}(m),$$

and m and M are, respectively, the least and greatest double zone diagrams. If $|K| = 2$, then by letting $m = (m_1, m_2)$ and $M = (M_1, M_2)$, each of the pairs (m_1, M_2) and (M_1, m_2) is a zone diagram.

Corollary 4.2. *Under the setting of Theorem 4.1, if X is compact, then by letting $M = (M_k)_{k\in K}$ and $m = (m_k)_{k\in K}$ we have*

$$M_k = \lim_{\gamma\to\infty} (\mathrm{Dom}^{2\gamma+1}(P))_k \quad \text{and} \quad m_k = \lim_{\gamma\to\infty} (\mathrm{Dom}^{2\gamma}(P))_k \quad \forall k \in K \tag{4}$$

with respect to the Hausdorff distance.

Corollary 4.3. *Under the setting of Theorem 4.1, if it is known that the least and the greatest double zone diagrams coincide, then $m = M$ and they both coincide with the unique zone diagram (even if X is not compact). In particular this is true when the normed space is Euclidean.*

Proof. By [22, Corollary 6.2] (whose proof is elementary) if the least and the greatest double zone diagram coincide, then, without any restriction on the sites or the space, there exists a unique zone diagram which is equal to both of them. Since in the proof of [12, Theorem 1.1] it was proved that the least double zone diagram coincides with the greatest one when the normed space is Euclidean, the assertion follows. □

Example 4.4. An illustration of Theorem 4.1 for the case where $m \neq M$ is given in Figures 6-8. Here the sites are $P_1 = \{(0, 1)\}$, $P_2 = \{(0, -1)\}$ in a rectangle in $\mathbb{R}^2$, but the norm is $\|(x_1, x_2)\| = \delta\sqrt{\alpha^2|x_1|^2 + |x_2|^2} + (1 - \delta\alpha)|x_1| + (1 - \delta)|x_2|$, where $\alpha = \delta = 0.1$. This norm is strictly convex but not smooth (the unit sphere contains points which have several supporting lines). A closely related example was discussed in [12, Section 5] in the context of non-uniqueness of zone diagrams (double zone diagrams were not mentioned) and it actually follows from the discussion presented there that indeed $m \neq M$ in this case. Two different zone diagrams can be obtained now, by taking either the first component of M together with the second of m or the second component of M together with the first of m.

For producing the pictures (see also Section 5) the endpoints of 4000 rays emanating from each site were computed in the Voronoi algorithm stage. However, it can be seen that still some parts of the boundaries of M are not full. The reason is that many rays should be produced in a very small angle (i.e., a very small part of the unit sphere) because of the location of the site with respect to these boundary parts. A simple way to overcome the problem (after detected) is to distribute the rays in the unit sphere in a non-uniform way such that many rays will emanate in the corresponding angle but relatively few in other angles, in contrast with the current way of producing the rays in a uniform way. One can also connect neighbor endpoints by line segments.

The proof of Theorem 4.1 is quite long and technical and it can be found in Section 6 together with the proof of Corollary 4.2 which follows from a quite general argument not



FIGURE 5. The components of the greatest double zone diagram M from Example 4.4.



FIGURE 6. The components of the least double zone diagram m from Example 4.4.



FIGURE 7. $I^{(2)}$ and $O^{(2)}$ for the double zone diagrams from Example 4.4.



FIGURE 8. $I^{(3)}$ and $O^{(3)}$ for the double zone diagrams from Example 4.4.

actually related to zone diagrams. Some of the ideas used in the proof of Theorem 4.1 are partly inspired from the ones given in [3, Lemma 3.1, Lemma 5.1]. However, the generalization of the arguments given there to our setting is definitely not immediate and one has to pay attention to certain subtle points, among them the verification of the equality

$$\bigcup_{j\neq k}\bigcap_{\gamma=1}^{\infty}(\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K})_j=\bigcap_{\gamma=1}^{\infty}\bigcup_{j\neq k}(\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K})_j$$

which is based on some non-immediate properties of dom and Dom.

It is interesting to note that Theorem 4.1 is also related to [10, Proposition 5] in the sense that in both cases one considers an iterative algorithm and the setting is closed sites in a finite dimensional strictly convex space. The proofs of these two claims have some similarities: for instance, [10, Lemma 6] is essentially Proposition 6.5 in Section 6. However, the proofs of these two claims are different in the strategy and the level of detailing (and historically they have been established independently). In general the proofs of [10, Proposition 5] and Theorem 4.1 here are different since the involved mappings are not the same and hence one needs to find a strategy which is appropriate for each case separately. In fact, an important feature used in the proof of Theorem 4.1 is the fact that the sites are positively separated (this also allows us to consider infinitely many sites) while in [10, Proposition 5] the sites satisfy the more general condition of being merely disjoint, but now one must consider only two sites (by definition) and the mapping must have finitely many components.

5. THE ACTUAL COMPUTATION OF $I^{(n)}$ AND $O^{(n)}$, TIME COMPLEXITY, AND CLARIFICATIONS

In this section we provide details about practical and theoretical aspects of the method which allows the approximate computation of the corresponding regions. The section has three part: in the first part practical details regarding the method are given. In the second part the issue of time complexity is considered. In the last part several related issues are clarified.

5.1. **The actual computation of $I^{(n)}$ and $O^{(n)}$:** As mentioned in Section 3, in order to compute the iterations $I^{(n)}$ and $O^{(n)}$ one has to know how to compute or approximate dominance regions $\text{dom}(P,A)$ of general sets P,A. The way we choose to overcome this difficulty is to use the approximation algorithm for computing Voronoi diagrams of general sites in general normed spaces which was introduced in [19]. In a nutshell, for approximating $\text{dom}(P,A)$ one uses the fact that $\text{dom}(P,A)$ can be represented as a union of line segments emanating from the points of P (see Figures 9-10 for an illustration).

First, the world X is assumed to be a large compact subset, e.g., a square or a hypercube. Now one approximates P using a finite collection of points; this is always possible if P is compact. After that a finite collection of directions (unit vectors) is chosen, and one approximates the endpoints of the line segments emanating from the points of P in these directions. These endpoints can be approximated to any required precision, and then they are stored. At this point $\text{dom}(P,A)$ is represented by this collection of endpoints and it is regarded as computed.

For computing $I^{(n)}$ and $O^{(n)}$ one computes their corresponding components iteratively: the components are $\text{dom}(P_k,A_k)$ where k runs over all the (finitely many in practice) indices in K and A_k depends on the iteration and on k. In general, $A_k=\bigcup_{j\neq k}(I^{(n)})_k$ or $A_k=\bigcup_{j\neq k}(O^{(n)})_k$ where $(I^{(n)})_k$ is the k-th component of $I^{(n)}$. For instance, the components of $I^{(1)}$ are $\text{dom}(P_k,A_k)$ where $A_k=\bigcup_{j\neq k}(O^{(0)})_k$ and $(O^{(0)})_k=\text{dom}(P_k,\bigcup_{j\neq k}P_j)$ is known, as a collection of endpoints, from a previous computation. An illustration of the process is given in Figures 3,4,11, and 12.

An important parameter which determines the level of approximation is the number of line segments (or, actually, the density of the corresponding unit vectors in the unit sphere of the space) used in the construction of the dominance regions. In Figures 3 and 4 about 1600 line segments emanate from each site, and only the endpoints are displayed. In Figures 8-fig:ZD-Iteration2-StrangeNorm-2Sites-0002 4000 rays emanate



FIGURE 9. The Voronoi diagram of 4 sites in $(\mathbb{R}^2, \ell_p)$, $p = 2.718281828$, each consists of 3 points; 80 line segments emanate from each point of each site.



FIGURE 10. The zone diagram (and hence the double zone diagram) of the sites of Figure 9.

from each site and only endpoints are displayed. In Figures 9-10 about 80 line segments are used, and the whole line segment is displayed.

5.2. **Time complexity:** As for the time complexity of the algorithm, it is determined by the time complexity of $\mathrm{dom}(P_k, A_k)$ for each k which is a particular case of the algorithm for computing Voronoi diagrams described in [19, 20]. Starting from iteration $n = 1$, the number of endpoints in A_k is $C_1 C_2^{d-1} G(|K| - 1)$ where C_1 is a universal positive constant depending on the norm, C_2 is a positive constant depending on the level of approximation of the unit sphere of the space (roughly speaking, it is a bound on the distance between a point on the unit sphere and some unit vector from the list of unit vectors created by the user), d is the dimension, G is a bound on the number of points in P_k (for each k), $C_1 C_2^{d-1} G$ is the number of line segments emanating from the points of P_k, and $|K|$ is the number of sites. For determining an endpoint up to some error parameter one makes $C_3 |A_k|$ comparisons where C_3 is a positive constant depending on the level of approximation. Hence the time complexity for the entire number of components is $O(C_3(C_1 C_2^{d-1} G|K|)^2)$. Note that here G and $|K|$ are parameters depending on the input, C_1 and d are global parameters, and C_2 and C_3 are parameters depending on the user (the level of approximation of the computation). Under some assumptions, e.g., that the sites are points which are uniformly distributed, it is possible to reduce significantly the number of calculations.

5.3. **Clarifications:** We finish this section with a discussion on certain issues which perhaps have not been so clear so far. It should be emphasized that the goal of this section is to describe schematically a practical way for approximating the regions which appear in each iteration and to evaluate roughly the number of calculations. The fact that the algorithm can approximate a given dominance region up to any desired precision is a simple consequence of the stability of the algorithm mentioned in [19].

However, the proof of this stability property is not obvious and it will be discussed elsewhere in a paper devoted to this algorithm. Its proof is in the spirit of the proof of the geometric stability of Voronoi diagrams with respect to small changes of the sites [21].

Roughly speaking, if the users want to approximate (in terms of the Hausdorff distance) a given region up to some error parameter ϵ, then they need to approximate well enough the sites, to choose in advance enough approximating rays (this is determined by the error parameter related to the unit sphere), and to fix a small enough error parameter for the endpoints of the rays. If the users decide in advance how many iterations they want to perform for approximating the (double) zone diagram and what is the level of approximation of the regions in the final iteration, then they can estimate in advance the number of calculations by iterative "reverse engineering": using the target error parameter ϵ, one estimates the error parameters needed as input for the final iteration, and from them the error parameters needed for the previous iteration, and so on, until the initial iteration. This gives an estimate on the initial error parameters.

In the above description the number of iterations n was chosen by the user but there was no guarantee that the real (double) zone diagram will be approximated well by $I^{(n)}$ or $O^{(n)}$. However, since the algorithm converges to the double zone diagram by Theorem 4.1, then given $\epsilon > 0$, there is a number n_0, depending only on ϵ, such that for any $n \geq n_0$, the regions of any iteration $I^{(n)}$ and $O^{(n)}$ will be at Hausdorff distance of at most ϵ from the limit regions. Hence, if one can compute $n_0(\epsilon)$, then one can know in advance the corresponding needed initial error parameters. Unfortunately, it is not clear how to estimate $n_0(\epsilon)$ and this is a major open problem.

6. PROOF OF THE MAIN RESULT

In this section we present the proof of Theorem 4.1 and the corollaries following it. We start with several lemmas describing simple properties of dom and Dom.

Lemma 6.1. *Let (X, d) be a metric space. Then*

(a) $d(x, \bigcup_{\gamma\in\Gamma} A_\gamma) = \inf\{d(x, A_\gamma) : \gamma \in \Gamma\}$ *for any $x \in X$ and any collection $\{A_\gamma\}_{\gamma\in\Gamma}$ of nonempty subsets in X.*

(b) $\mathrm{dom}(P, \bigcup_{\gamma\in\Gamma} A_\gamma) = \bigcap_{\gamma\in\Gamma} \mathrm{dom}(P, A_\gamma)$ *for any collection $\{A_\gamma\}_{\gamma\in\Gamma}$ of nonempty subsets in X and any $P \subseteq X$ nonempty.*

(c) $\mathrm{dom}(\bigcup_{\gamma\in\Gamma} P_\gamma, A) \subseteq \bigcup_{\gamma\in\Gamma} \mathrm{dom}(P_\gamma, A)$ *for any collection $\{P_\gamma\}_{\gamma\in\Gamma}$ of nonempty subsets in X and any $A \subseteq X$ nonempty. If, in addition, for each $x \in X$ the distance $d(x, \bigcup_{\gamma\in\Gamma} P_\gamma)$ is attained at some P_{γ_0}, i.e., $d(x, \bigcup_{\gamma\in\Gamma} P_\gamma) = d(x, P_{\gamma_0})$ for some $\gamma_0 \in \Gamma$, then the equality* $\mathrm{dom}(\bigcup_{\gamma\in\Gamma} P_\gamma, A) = \bigcup_{\gamma\in\Gamma} \mathrm{dom}(P_\gamma, A)$ *holds.*

(d) $\mathrm{Dom}(\bigcup_{\gamma\in\Gamma} R^\gamma) = \bigcap_{\gamma\in\Gamma} \mathrm{Dom}(R^\gamma)$ *for any collection $\{R^\gamma\}_{\gamma\in\Gamma}$ of tuples, where each R^γ is indexed by the same set of indices K, i.e., $R^\gamma = (R^\gamma_k)_{k\in K}$. Here* Dom *is defined with respect to some tuple $(P_k)_{k\in K}$ of nonempty subsets in X.*

Proof. (a) Let $x \in X$, $\alpha := d(x, \bigcup_{\gamma\in\Gamma} A_\gamma)$ and $\beta := \inf\{d(x, A_\gamma) : \gamma \in \Gamma\}$. Then $\alpha \leq d(x, A_\gamma)$ for all $\gamma \in \Gamma$ by the definition of α, so $\alpha \leq \beta$. If $\alpha < \beta$, then there is $y \in \bigcup_{\gamma\in\Gamma} A_\gamma$ such that $d(x, y) < \beta$, and since $y \in A_\gamma$ for some $\gamma \in \Gamma$, we have $d(x, A_\gamma) \leq d(x, y) < \beta$, a contradiction with the definition of β.

(b) By the antimonotonicity of $\mathrm{dom}(P,\cdot)$ we have $\mathrm{dom}(P,\bigcup_{i\in\Gamma}A_i)\subseteq\mathrm{dom}(P,A_\gamma)$ for any $\gamma\in\Gamma$. Consequently, we have $\mathrm{dom}(P,\bigcup_{\gamma\in\Gamma}A_\gamma)\subseteq\bigcap_{\gamma\in\Gamma}\mathrm{dom}(P,A_\gamma)$. Conversely, suppose that $x\in\bigcap_{\gamma\in\Gamma}\mathrm{dom}(P,A_\gamma)$. If $x\notin\mathrm{dom}(P,\bigcup_{\gamma\in\Gamma}A_\gamma)$, then there is $y\in\bigcup_{\gamma\in\Gamma}A_\gamma$ such that $d(x,P)>d(x,y)$. But $y\in A_\gamma$ for some γ, and $x\in\mathrm{dom}(P,A_\gamma)$, so $d(x,P)\le d(x,A_\gamma)\le d(x,y)$, a contradiction.

(c) By monotonicity of $\mathrm{dom}(\cdot,A)$, i.e., $\mathrm{dom}(P,A)\le\mathrm{dom}(Q,A)$ whenever $P\subseteq Q$ (which is easily verified), we have $\mathrm{dom}(P_\gamma,A)\subseteq\mathrm{dom}(\bigcup_{i\in\Gamma}P_i,A)$ for any γ. As a result, we have $\bigcup_{\gamma\in\Gamma}\mathrm{dom}(P_\gamma,A)\subseteq\mathrm{dom}(\bigcup_{\gamma\in\Gamma}P_\gamma,A)$. Now suppose that $d(x,\bigcup_{\gamma\in\Gamma}P_\gamma)=d(x,P_{\gamma_0})$ for some $\gamma_0\in\Gamma$. Therefore, if $x\in\mathrm{dom}(\bigcup_{\gamma\in\Gamma}P_\gamma,A)$, then $d(x,P_{\gamma_0})=d(x,\bigcup_{\gamma\in\Gamma}P_\gamma)\le d(x,A)$, i.e., $x\in\mathrm{dom}(P_{\gamma_0},A)\subseteq\bigcup_{\gamma\in\Gamma}\mathrm{dom}(P_\gamma,A)$.

(d) Since the union and intersection are taken component-wise, this follows from the definition of the Dom mapping (equation (1)) and part (b):

$$\begin{aligned}\mathrm{Dom}(\bigcup_{\gamma\in\Gamma}R^\gamma)=(\mathrm{dom}(P_k,\bigcup_{j\neq k}(\bigcup_{\gamma\in\Gamma}R_j^\gamma)))_{k\in K}&=(\mathrm{dom}(P_k,\bigcup_{\gamma\in\Gamma}(\bigcup_{j\neq k}R_j^\gamma)))_{k\in K}\\&=(\bigcap_{\gamma\in\Gamma}\mathrm{dom}(P_k,\bigcup_{j\neq k}R_j^\gamma))_{k\in K}=\bigcap_{\gamma\in\Gamma}\mathrm{Dom}(R^\gamma).\end{aligned}$$

□

Lemma 6.2. *Let $(P_k)_{k\in K}$ be a tuple of nonempty and closed subsets in X.*

(a) Dom *is antimonotone, i.e.,* $\mathrm{Dom}(R)\subseteq\mathrm{Dom}(S)$ *whenever* $S\subseteq R$*;* Dom^2 *is monotone, that is,* $R\subseteq S\Rightarrow\mathrm{Dom}^2(R)\subseteq\mathrm{Dom}^2(S)$.

(b) The sequence $(I^{(n)})_{n=0}^\infty$ is increasing, $(O^{(n)})_{n=0}^\infty$ is decreasing and we have

$$I^{(n)}\subseteq\bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P_k)_{k\in K}\subseteq\bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}\subseteq O^{(n)}\tag{5}$$

for each nonnegative n. In addition, if R is a zone or double zone diagram in X, then

$$I^{(n)}\subseteq\bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P_k)_{k\in K}\subseteq R\subseteq\bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}\subseteq O^{(n)}\quad\forall n\ge 0.\tag{6}$$

Proof. (a) The antimonotonicity of Dom is a simple consequence of the easily verified fact that $\mathrm{dom}(P,B)\subseteq\mathrm{dom}(P,A)$ for any nonempty subset P and any nonempty subsets A,B satisfying $A\subseteq B$. The composition of two antimonotone mapping is monotone, and hence Dom^2 is monotone.

(b) Let $(X)_{k\in K}$ be the tuple whose components are X. An immediate check shows that $P_k=\mathrm{dom}(P_k,X)$ and $\mathrm{Dom}(X)_{k\in K}=(P_k)_{k\in K}$. From the inclusion $(P_k)_{k\in K}\subseteq(X)_{k\in K}$, the definition of the space on which Dom acts, and the monotonicity of Dom^2 we have

$$(P_k)_{k\in K}\subseteq\mathrm{Dom}^2(P_k)_{k\in K}\subseteq\mathrm{Dom}^2(X)_{k\in K}=\mathrm{Dom}(P_k)_{k\in K}\subseteq(X)_{k\in K}.$$

As a result, by iterating with Dom^2, using its monotonicity, and using the definition of $I^{(n)}$ and $O^{(n)}$ we see that $(I^{(n)})_{n=0}^\infty$ is increasing and $(O^{(n)})_{n=0}^\infty$ is decreasing. Hence $I^{(n)}$ is contained in $\bigcup_{\ell=0}I^{(\ell)}$ and $\bigcap_{\ell=0}O^{(\ell)}$ is contained in $O^{(n)}$ for each n. Finally, after iterating the inclusion $(P_k)_{k\in K}\subseteq R\subseteq(X)_{k\in K}$ using Dom^2 we obtain (6) when R is a double zone diagram. But a zone diagram is a double zone diagram

[$\mathrm{Dom}(\mathrm{Dom}(R))=\mathrm{Dom}(R)=R$ whenever $R=\mathrm{Dom}(R)$], so (6) is true in this case too.

□

Lemma 6.3. *Let (X,d) be a metric space and let $P,A\subseteq X$ be nonempty. Then*

(a) $\mathrm{dom}(P,A)$ *is a closed set.*

(b) $\mathrm{dom}(P,A)=\mathrm{dom}(P,\overline{A})=\mathrm{dom}(\overline{P},A)=\mathrm{dom}(\overline{P},\overline{A})$.

(c) For a tuple $R=(R_k)_{k\in K}$ of nonempty subsets, let $\overline{R}=(\overline{R_k})_{k\in K}$. Then $\mathrm{Dom}(\overline{R})=\mathrm{Dom}(R)=\overline{\mathrm{Dom}(R)}$ *(with respect to a given tuple $(P_k)_{k\in K}$).*

Proof. (a) If $x=\lim_{n\to\infty}x_n$ and $x_n\in\mathrm{dom}(P,A)$, then $d(x_n,P)\le d(x_n,A)$, and this inequality is preserved in the limit because the function $x\mapsto d(x,P)-d(x,A)$ is continuous (with respect to the topology induced on X by d).

(b) This follows from the general fact that $d(x,B)=d(x,\overline{B})$ for each $x\in X$ and each $B\subseteq X$.

(c) By part (b) and by Lemma 6.1(b), we have

$$\mathrm{dom}(P_k,\bigcup_{j\neq k}\overline{R_j})=\bigcap_{j\neq k}\mathrm{dom}(P_k,\overline{R_j})=\bigcap_{j\neq k}\mathrm{dom}(P_k,R_j)=\mathrm{dom}(P_k,\bigcup_{j\neq k}R_j).$$

Thus $\mathrm{Dom}(\overline{R})=(\mathrm{dom}(P_k,\bigcup_{j\neq k}\overline{R_j}))_{k\in K}=(\mathrm{dom}(P_k,\bigcup_{j\neq k}R_j))_{k\in K}=\mathrm{Dom}(R)$. Finally, by part (a) and by definition, we have

$$\overline{\mathrm{Dom}(R)}=(\overline{\mathrm{dom}(P_k,\bigcup_{j\neq k}P_j)})_{k\in K}=(\mathrm{dom}(P_k,\bigcup_{j\neq k}P_j))_{k\in K}=\mathrm{Dom}(R).$$

□

Proposition 6.4. *Let X be a convex subset of a finite dimensional strictly convex normed space $(\widetilde{X},|\cdot|)$. Let $A\subseteq X$ be nonempty and closed. Let $p,z\in X$ and suppose that $d(z,p)\le d(z,A)$ and $p\notin A$. Then $d(x,p)<d(x,A)$ for all $x\in[p,z)$.*

Proof. We can assume that $z\neq p$, because otherwise the assertion is obvious (void). Fix $x\in[p,z)$ and let $a_x\in A$ be chosen such that $d(x,A)=d(x,a_x)$. There exists such an a_x by compactness since the space is finite dimensional and A is closed. Since $p\notin A$ and $d(z,p)\le d(z,A)$ it follows that z,p and a_x are all different.

We distinguish between several cases. Suppose first that p,z,a_x are on the same line. If p is between z and a_x, then obviously $d(x,p)<d(x,a_x)=d(x,A)$. The case where a_x is between z and p is impossible, since then $d(z,a_x)<d(z,p)$, contradicting $d(z,p)\le d(z,A)$. If z is between p and a_x, then $d(z,a_x)<d(z,x)+d(x,a_x)$. This inequality is true also if p,z and a_x are not on the same line, by the strict convexity of the norm. Hence, in these two latter cases we have

$$d(z,x)+d(x,p)=d(z,p)\le d(z,A)\le d(z,a_x)<d(z,x)+d(x,a_x),$$

so $d(x,p)<d(x,a_x)=d(x,A)$. □

Proposition 6.5. *Let X be a convex subset of a finite dimensional strictly convex normed space $(\widetilde{X},|\cdot|)$. Let $P,A\subseteq X$ be nonempty and closed, and suppose that $P\bigcap A=\emptyset$. Then*

$$\mathrm{dom}(P,A)\ =\ \overline{\{x\in X: d(x,P)<d(x,A)\}}.$$

Proof. From the inclusion $\{x \in X : d(x,P) < d(x,A)\} \subseteq \mathrm{dom}(P,A)$ and the fact that $\mathrm{dom}(P,A)$ is closed (Lemma 6.3(a)) we have the inclusion $\overline{\{x \in X : d(x,P) < d(x,A)\}} \subseteq \mathrm{dom}(P,A)$. For the second inclusion, let $z \in \mathrm{dom}(P,A)$. If $d(z,P) < d(z,A)$, then obviously the point z is in the set $\overline{\{x \in X : d(x,P) < d(x,A)\}}$. Now suppose that $d(z,P) = d(z,A)$. By our assumption there is $p \in P$ such that $d(z,P) = d(z,p)$, and since $p \in P$, it follows that $p \notin A$. Since A is closed we have $d(p,A) > 0$. Hence if $z = p$, then $z \in \overline{\{x \in X : d(x,P) < d(x,A)\}}$. If $z \neq p$, then $[p,z) \neq \emptyset$, and Proposition 6.4 implies that any $x \in [p,z)$ (arbitrary close to z) satisfies the inequality $d(x,P) \leq d(x,p) < d(x,A)$. Thus again $z \in \overline{\{x \in X : d(x,P) < d(x,A)\}}$.

□

The following lemmas are somewhat independent of the main concepts of the paper but they are needed later.

Lemma 6.6. *Let X be a closed subset in a normed space $(\widetilde{X}, |\cdot|)$ and let $(A_\gamma)_{\gamma=1}^{\infty}$ be a decreasing sequence of subsets of X such that $A := \bigcap_{\gamma=1}^{\infty} A_\gamma$ is nonempty. If A_γ is closed for each γ and $(\widetilde{X}, |\cdot|)$ is finite dimensional or X is compact, then $d(x,A) = \lim_{\gamma\to\infty} d(x,A_\gamma)$ for each $x \in X$.*

Proof. Let $x \in X$ and assume that $(\widetilde{X}, |\cdot|)$ is finite dimensional or X is compact. In this case compactness arguments yield that the distance between x and any nonempty closed subset is attained. Therefore for each γ there exists $x_\gamma \in A_\gamma$ such that $d(x,A_\gamma) = d(x,x_\gamma)$. Since $(A_\gamma)_{\gamma=1}^{\infty}$ is decreasing and A_γ is closed for each γ, it follows that any limit point of the sequence $(x_\gamma)_{\gamma\in\mathbb{N}}$ is in A_γ for each γ and hence in A. Since $(A_\gamma)_{\gamma=1}^{\infty}$ is decreasing, the sequence $(d(x,x_\gamma))_{\gamma\in\mathbb{N}}$ is increasing and its limit is bounded by $d(x,A)$. Hence $(x_\gamma)_{\gamma\in\mathbb{N}}$ is bounded and there exists $x_\infty \in A$ and a subsequence such that $x_\infty = \lim_{\beta\to\infty} x_{\gamma_\beta}$. From the continuity of the distance function we deduce that $d(x,x_\infty) = \lim_{\beta\to\infty} d(x,x_{\gamma_\beta}) \leq d(x,A)$. But $d(x,A) \leq d(x,x_\infty)$ by definition, so $\lim_{\gamma\to\infty} d(x,A_\gamma) = \lim_{\beta\to\infty} d(x,A_{\gamma_\beta}) = d(x,x_\infty) = d(x,A)$. □

Lemma 6.7. *Let (X,d) be a compact metric space. Let $(A_n)_{n=1}^{\infty}$ be a decreasing sequence of nonempty closed sets of X, and let $(B_n)_{n=1}^{\infty}$ be an increasing sequence of nonempty sets of X. Let $A = \bigcap_{n=1}^{\infty} A_n$ and $B = \overline{\bigcup_{n=1}^{\infty} B_n}$. Then $A = \lim_{n\to\infty} A_n$ and $B = \lim_{n\to\infty} B_n$ with respect to the Hausdorff distance.*

Proof. Suppose by way of contradiction that it is not true that $\lim_{n\to\infty} D(A_n,A) = 0$, where D is the Hausdorff distance. Then there exists some $\epsilon > 0$ and a subsequence $n_m \in \mathbb{N}$ such that $\epsilon \leq D(A_{n_m},A)$ for each $m \in \mathbb{N}$. Since $A \subseteq A_n$ for each n and because X is compact, the definition of the Hausdorff distance implies that for each m there exists $x_m \in A_{n_m}$ such that $\epsilon \leq d(x_m,A)$. Let $(x_{m_l})_{l\in\mathbb{N}}$ be a convergent subsequence of the sequence $(x_m)_{m=1}^{\infty}$, and let x be its limit. Since the sequence $(A_n)_n$ is decreasing, it follows that $x_{m_{l'}} \in A_{n_{m_l}}$ whenever $l \leq l'$. Therefore $x \in A_{n_{m_l}}$ because $A_{n_{m_l}}$ is closed. Thus $x \in \bigcap_{l=1}^{\infty} A_{n_{m_l}}$. But $\bigcap_{l=1}^{\infty} A_{n_{m_l}} = A$ because the intersection is decreasing. Hence $0 = d(x,A) = \lim_{l\to\infty} d(x_{m_l},A) \geq \epsilon$, a contradiction.

Now suppose by way of contradiction that it is not true that $\lim_{n\to\infty} D(B_n,B) = 0$. Then there exists some $\epsilon > 0$ and a subsequence $n_m \in \mathbb{N}$ such that $\epsilon \leq D(B_{n_m},B)$ for each $m \in \mathbb{N}$. Since $B_n \subseteq B$ for each n and X is compact, the definition of the Hausdorff distance implies that for each m there exists $x_m \in B$ such that $\epsilon \leq d(x_m,B_{n_m})$. Let

$(x_{m_l})_{l\in\mathbb{N}}$ be a convergent subsequence of the sequence $(x_m)_{m=1}^{\infty}$, and let x be its limit. Then $d(x,x_{m_l})<\epsilon/4$ for all l large enough. Since B is closed it follows that $x\in B$, so by the definition of B there exists $n_0\in\mathbb{N}$ such that $d(x,B_{n_0})<\epsilon/2$. But $(B_n)_{n=1}^{\infty}$ is increasing, so $d(x,B_n)<\epsilon/2$ for all $n\in\mathbb{N}$ such that $n\geq n_0$, and, in particular, $d(x,B_{n_{m_l}})<\epsilon/2$ for l large enough. Thus

$$\epsilon\leq d(x_{m_l},B_{n_{m_l}})\leq d(x_{m_l},x)+d(x,B_{n_{m_l}})<3\epsilon/4,$$

a contradiction. □

In the next claims more subtle properties of dom and Dom will be derived. Lemma 6.10 and Proposition 6.11 generalize [3, Lemma 3.1] and [3, Lemma 5.1] respectively, and are partly inspired by them.

Lemma 6.8. *Let (X,d) be a metric space and let $P=(P_k)_{k\in K}$ be a tuple of nonempty subsets of X such that for all $k\in K$,*

$$r_k:=\inf\{d(P_k,P_j):j\neq k\}>0.$$

For $Q\subseteq X$ nonempty and $r>0$, let $B(Q,r)=\{x\in X:d(x,Q)<r\}$. We denote by $(\mathrm{Dom}P)_k$ the k-th component of $\mathrm{Dom}P$. Then:

(a) $B(P_k,r_k/4)\subseteq(\mathrm{Dom}^2P)_k\subseteq(\mathrm{Dom}^{\gamma}P)_k$ for any integer $\gamma\geq 1$ and any $k\in K$.

(b) If X is a convex subset of a normed space, then $d(x,B(Q,r))+r\leq d(x,Q)$ for any $Q\subseteq X$ nonempty, $r>0$ and $x\notin B(Q,r)$.

(c) If $d(x,B(P_k,r))<d(x,P_k)$ for any $k\in K$, $r>0$, and $x\notin\overline{P_k}$, then for each integer $\gamma\geq 2$ the components of $\mathrm{Dom}^{\gamma}P$ are disjoint. Moreover, if X is a convex subset of a normed space, then for any $j,k\in K, j\neq k$ and any $\gamma\geq 2$ we have $r_k/8+r_j/8\leq d((\mathrm{Dom}^{\gamma}P)_k,(\mathrm{Dom}^{\gamma}P)_j)$.

(d) Suppose that X is a convex subset of a normed space. Given $j,k\in K$, $j\neq k$, and an integer $\gamma\geq 2$, if $x\in(\mathrm{Dom}^{\gamma}P)_j$, then $r_k/4\leq d(x,P_k)$.

Proof. (a) Let $k\in K$ and suppose that $x\in B(P_k,r_k/4)$, i.e., $d(x,P_k)<r_k/4$. By definition we have $(\mathrm{Dom}^2P)_k=\mathrm{dom}(P_k,\bigcup_{j\neq k}\mathrm{dom}(P_j,\bigcup_{i\neq j}P_i))$, so in order to prove that $x\in(\mathrm{Dom}^2P)_k$ it suffices to prove that $r_k/4\leq d(x,y)$ for all $y\in\bigcup_{j\neq k}\mathrm{dom}(P_j,\bigcup_{i\neq j}P_i)$. Given y in this union, it is in $\mathrm{dom}(P_j,\bigcup_{i\neq j}P_i)$ for some $j\neq k$. Hence

$$d(y,P_j)\leq d(y,\bigcup_{i\neq j}P_i)\leq d(y,P_k)\leq d(y,x)+d(x,P_k).$$

Therefore, by the definition of r_k,

$$r_k\leq d(P_j,P_k)\leq d(P_j,y)+d(y,x)+d(x,P_k)\leq 2d(x,y)+2r_k/4,$$

i.e., $d(x,P_k)<r_k/4\leq d(x,y)$ and the assertion follows. Finally, Lemma 6.2 implies that $\mathrm{Dom}^2P\subseteq\mathrm{Dom}^{\gamma}P$ for any integer $\gamma\geq 1$, and hence $(\mathrm{Dom}^2P)_k\subseteq(\mathrm{Dom}^{\gamma}P)_k$ for each $k\in K$.

(b) Let $p\in Q$. Since $x\notin B(Q,r)$, a simple argument shows that the intersection of the compact segment $[x,p]$ with the closed set $\partial B(Q,r)$ is a nonempty compact set, and hence the distance between x and this intersection is attained at some point $y\in[x,p]$. It must be that $s:=r-d(y,p)\leq 0$. If this is not true, then for any z in the open ball $B(y,s)$ we have $d(z,Q)\leq d(z,p)\leq d(z,y)+d(y,p)<$

$s+d(y,p)=r$, and hence y is in the interior of $B(Q,r)$, a contradiction. Thus $d(x,p)=d(x,y)+d(y,p)\geq d(x,y)+r$, so

$$d(x,B(Q,r))=d(x,\overline{B(Q,r)})\leq d(x,y)\leq d(x,p)-r,$$

for any $p\in Q$. As a result, $d(x,B(Q,r))+r\leq d(x,Q)$.

(c) Suppose that we know that the components of Dom^3P are disjoint. Then for each integer $\gamma\geq 2$ the components of $\mathrm{Dom}^{\gamma}P$ are disjoint, because $\mathrm{Dom}^{\gamma}P\subseteq\mathrm{Dom}^3P$ by Lemma 6.2. We now give a proof for the case $\gamma=3$. Let $k_1\neq k_2$ be two indices in K, and assume to the contrary that $x\in(\mathrm{Dom}^3P)_{k_1}\bigcap(\mathrm{Dom}^3P)_{k_2}$. By definition and by part (a) we have

$$\begin{aligned}d(x,P_{k_1})\leq d(x,\bigcup_{j\neq k_1}(\mathrm{Dom}^2P)_j)&\leq d(x,(\mathrm{Dom}^2P)_{k_2})\\&\leq d(x,B(P_{k_2},r_{k_2}/4))\leq d(x,P_{k_2}).\end{aligned}$$

Hence $x\notin\overline{P_{k_2}}$, because otherwise also $x\in\overline{P_{k_1}}$, a contradiction with the assumption that $0<r_{k_1}\leq d(P_{k_1},P_{k_2})$. Thus $d(x,B(P_{k_2},r_{k_2}/4))<d(x,P_{k_2})$ by assumption, so actually $d(x,P_{k_1})<d(x,P_{k_2})$. In the same way $d(x,P_{k_2})<d(x,P_{k_1})$, a contradiction.

Finally, suppose that X is a convex subset of a normed space. Let $j,k\in K$ be different. Since $(\mathrm{Dom}^{\gamma}P)_i\subseteq(\mathrm{Dom}^3P)_i$ for each integer $\gamma\geq 2$ and each $i\in K$, it suffices to show that $r_k/8+r_j/8\leq d(x,y)$ for any $x\in(\mathrm{Dom}^3P)_k$ and $y\in(\mathrm{Dom}^3P)_j$. By definition, the triangle inequality and parts (a),(b),

$$\begin{aligned}d(x,P_k)\leq d(x,\bigcup_{i\neq k}(\mathrm{Dom}^2P)_i)\leq d(x,(\mathrm{Dom}^2P)_j)&\leq d(x,B(P_j,r_j/4))\\&\leq d(x,P_j)-r_j/4\leq d(x,y)+d(y,P_j)-r_j/4.\end{aligned}$$

In the same way $d(y,P_j)\leq d(y,x)+d(x,P_k)-r_k/4$. By adding these two inequalities we obtain the desired conclusion.

(d) If $x\in(\mathrm{Dom}^{\gamma}(P))_j$, then $x\notin B(P_k,r_k/4)\subseteq(\mathrm{Dom}^{\gamma}(P))_k$ by parts (a) and (c). Hence $r_k/4\leq d(x,P_k)$ by part (b).

□

Remark 6.9. Lemma 6.8 parts (b),(c),(d) hold with exactly the same proof in any geodesic metric space. Recall that a metric space (X,d) is called geodesic if between any two points x and y in it there exists an isometry γ (i.e., γ preserves distances) from the real line segment $[0,d(y,x)]$ to X such that $\gamma(0)=x$ and $\gamma(d(y,x))=y$. Simple and familiar examples of geodesic metric spaces are: the Euclidean plane, any convex subset of a normed space, spheres, complete Riemannian manifolds [11, pp. 25-28], and hyperbolic spaces [23, pp. 538-9].

Lemma 6.10. *Let (X,d) be a metric space, let $P\subseteq X$ be nonempty and suppose that $\{A_i\}_{i=1}^{\infty}$ is a family of subsets of X such that $\bigcap_{i=1}^{\infty}A_i\neq\emptyset$. If*

$$\mathrm{dom}(P,\bigcap_{i=1}^{\infty}A_i)=\overline{\{x\in X:d(x,P)<d(x,\bigcap_{i=1}^{\infty}A_i)\}},\tag{7}$$

and

$$d(y,\bigcap_{i=1}^{\infty}A_i)=\limsup_{i\to\infty}d(y,A_i),\quad \forall y\in X,\tag{8}$$

then

$$\mathrm{dom}(P,\bigcap_{i=1}^{\infty}A_i)=\overline{\bigcup_{i=1}^{\infty}\mathrm{dom}(P,A_i)}.\tag{9}$$

Proof. By antimonotonicity, $\mathrm{dom}(P,\bigcap_{j=1}^{\infty}A_j)\supseteq\mathrm{dom}(P,A_i)$ for all i, and since the left hand side is closed, the inclusion $\mathrm{dom}(P,\bigcap_{i=1}^{\infty}A_i)\supseteq\overline{\bigcup_{i=1}^{\infty}\mathrm{dom}(P,A_i)}$ follows. For the reverse inclusion, let $\epsilon>0$ be given and suppose that $x\in\mathrm{dom}(P,\bigcap_{i=1}^{\infty}A_i)$. We should prove that there are $i\in\mathbb{N}$ and $y\in\mathrm{dom}(P,A_i)$ such that $d(y,x)<\epsilon$.

By (7) there is $y\in X$ such that $d(x,y)<\epsilon$ and $r:=d(y,\bigcap_{i=1}^{\infty}A_i)-d(y,P)>0$, and by (8) there is i large enough such that $|d(y,\bigcap_{j=1}^{\infty}A_j)-d(y,A_i)|<r/2$. Hence

$$d(y,P)+r/2<d(y,\bigcap_{i=1}^{\infty}A_i)<d(y,A_i)+r/2.$$

Thus $d(y,P)<d(y,A_i)$, and so $y\in\mathrm{dom}(P,A_i)$. $\square$

Proposition 6.11. *Let X be a be a convex set in a finite dimensional strictly convex space $(\widetilde{X},|\cdot|)$ and let $(P_k)_{k\in K}$ be a tuple of nonempty closed sets in X such that $r:=\inf\{d(P_k,P_j):j,k\in K,\,j\neq k\}>0$. Then*

$$\mathrm{Dom}(\bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K})=\overline{\bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P_k)_{k\in K}}.$$

Proof. For each nonnegative integer γ, let $R^{\gamma}=\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}$. We claim that

$$\bigcup_{j\neq k}\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}=\bigcap_{\gamma=1}^{\infty}\bigcup_{j\neq k}R_j^{\gamma}.\tag{10}$$

Indeed, if $x\in\bigcup_{j\neq k}\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}$, then $x\in\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}$ for some $j\neq k$. Since $R_j^{\gamma}\subseteq\bigcup_{i\neq k}R_i^{\gamma}$, it follows that $\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}\subseteq\bigcap_{\gamma=1}^{\infty}\bigcup_{i\neq k}R_i^{\gamma}$, so $x\in\bigcap_{\gamma=1}^{\infty}\bigcup_{i\neq k}R_i^{\gamma}$. On the other hand, let $x\in\bigcap_{\gamma=1}^{\infty}\bigcup_{j\neq k}R_j^{\gamma}$. Given $\gamma\in\mathbb{N}$, Lemma 6.8(c) (note that $2\gamma+1\geq 3$ and $R^{\gamma}=\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}$) implies that $R_i^{\gamma}\bigcap R_j^{\gamma}=\emptyset$ for $j\neq i$, so there exists exactly one index $j\neq k$ such that $x\in R_j^{\gamma}$. It must be that all these indices coincide. If this is not true, then $x\in R_j^{\gamma}\bigcap R_{j'}^{\gamma'}$ for some $\gamma'>\gamma$ and corresponding indices $j'\neq j$. But $R_{j'}^{\gamma'}\subseteq R_{j'}^{\gamma}$ by Lemma 6.2, so $x\in R_j^{\gamma}\bigcap R_{j'}^{\gamma}$, a contradiction to Lemma 6.8(c). As a result $x\in\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}$ for some $j\neq k$, and this establishes (10).

Now fix $k\in K$ and let $A_{\gamma}=\bigcup_{j\neq k}R_j^{\gamma}$ and $A=\bigcap_{\gamma=1}^{\infty}A_{\gamma}$. By Lemma 6.2 we have $P_j\subseteq\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}$, so (10) implies that $A\neq\emptyset$. By Lemma 6.8(d) it follows that $0<r/4\leq d(P_k,A)$.

By Lemma 6.2 the intersection which defines A is decreasing. By Lemma 6.3(a) and Lemma 6.8(c) each A_{γ} is a closed set, because it is a union of closed and disjoint sets with a positive distance $r/4$ between any two different members in the union. Therefore Lemma 6.6 implies that $d(x,A)=\lim_{\gamma\to\infty}d(x,A_{\gamma})$ and hence (8) holds. In addition, from the above we see that A is closed and hence we conclude from Proposition 6.5

that (7) holds for $P = P_k$. Consequently, we conclude from Lemma 6.2, the definition of Dom, (10), and Lemma 6.10 that

$$\begin{aligned}
\mathrm{Dom}\left(\bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P_k)_{k\in K}\right) &= \mathrm{Dom}\left(\bigcap_{\gamma=0}^{\infty}R^{\gamma}\right) = \mathrm{Dom}\left(\bigcap_{\gamma=1}^{\infty}R^{\gamma}\right)\\
&= \left(\mathrm{dom}\left(P_k,\bigcup_{j\neq k}\bigcap_{\gamma=1}^{\infty}R_j^{\gamma}\right)\right)_{k\in K} = \left(\mathrm{dom}\left(P_k,\bigcap_{\gamma=1}^{\infty}\bigcup_{j\neq k}R_j^{\gamma}\right)\right)_{k\in K}\\
&= \left(\overline{\bigcup_{\gamma=1}^{\infty}\mathrm{dom}(P_k,\bigcup_{j\neq k}R_j^{\gamma})}\right)_{k\in K} = \overline{\bigcup_{\gamma=1}^{\infty}\mathrm{Dom}(R^{\gamma})} = \overline{\bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P_k)_{k\in K}}.
\end{aligned}$$

□

We are now able to prove the main result.

Proof of Theorem 4.1. Part of the proof is completely general. By Lemma 6.1(d) and Lemma 6.3(c) we obtain

$$\mathrm{Dom}(m) = \mathrm{Dom}\left(\overline{\bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P)}\right) = \bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P) = M. \tag{11}$$

On the other hand, Proposition 6.11 implies that

$$\mathrm{Dom}(M) = \mathrm{Dom}\left(\bigcap_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma+1}(P)\right) = \overline{\bigcup_{\gamma=0}^{\infty}\mathrm{Dom}^{2\gamma}(P)} = m. \tag{12}$$

Thus $m = \mathrm{Dom}(M) = \mathrm{Dom}(\mathrm{Dom}(m))$ and $M = \mathrm{Dom}^2(M)$, so both m and M are double zone diagrams. From Lemma 6.2 and Lemma 6.3(c) we see that m and M are respectively the least and greatest double zone diagrams. Finally, if $|K| = 2$, then by (11), (12), and the definition of Dom we have $(M_1, M_2) = \mathrm{Dom}(m) = ((\mathrm{dom}(P_1, m_2), \mathrm{dom}(P_2, m_1))$ and $(m_1, m_2) = (\mathrm{dom}(P_1, M_2), \mathrm{dom}(P_2, M_1))$. Hence $m_1 = \mathrm{dom}(P_1, M_2)$ and $M_2 = \mathrm{dom}(P_2, m_1)$. But the definition of Dom implies the equality $\mathrm{Dom}(m_1, M_2) = (\mathrm{dom}(P_1, M_2), \mathrm{dom}(P_2, m_1))$. Therefore $\mathrm{Dom}(m_1, M_2) = (m_1, M_2)$, i.e., (m_1, M_2) is a zone diagram. By the same way (M_1, m_2) is a zone diagram.

□

Proof of Corollary 4.2. By Lemma 6.2 the sequence $A_\gamma = (\mathrm{Dom}^{2\gamma+1}(P))_k$ is decreasing and the sequence $B_\gamma = (\mathrm{Dom}^{2\gamma}(P))_k$ is increasing for each $k \in K$. As a result, when X is compact (4) follows from Lemma 6.7. □

7. Open problems

We conclude this paper with several interesting open problems and unexplained phenomena. First, perhaps the most interesting open problem is to establish error estimates for the convergence speed of $(I^{(n)})_{n=0}^{\infty}$ and $(O^{(n)})_{n=0}^{\infty}$. One of the reasons that this problem is interesting, is the observed fast convergence of these sequences (usually 4 iterations suffice). We feel that the approach and error estimates developed in [21] may help here.



FIGURE 11. $I^{(1)}, \dots, I^{(4)}$ and $O^{(1)}, \dots, O^{(4)}$ for the sites $\{(0,3)\}$, $\{(0,-3)\}$ in a square in $(\mathbb{R}^2, \ell_\infty)$; the bold line is $I^{(3)} \approx I^{(4)} \approx O^{(3)} \approx O^{(4)}$. See also [22, Example 2.4].



FIGURE 12. $I^{(2)}$ and $O^{(2)}$ for 30 point sites in a square in $(\mathbb{R}^2, \ell_1)$.

Second, it is interesting to find out whether Theorem 4.1 can be generalized to other settings, for instance to all normed spaces, or at least to finite dimensional ones. Recently we have made some progress regarding this issue in the particular case of infinite dimensional uniformly convex spaces. This issue will be discussed in a future version.

Third, although in general a zone diagram is not necessarily unique, even in the case of $\mathbb{R}^2$ with two point sites (see e.g., [12, Sections 1,5] or Example 4.4), and although our results are limited to strictly convex spaces, experiments show that the algorithm actually converges to a unique zone diagram most of the times also in the case of non-strictly convex spaces; see Figures 11-12 for an illustration.

ACKNOWLEDGMENTS

To be added.

REFERENCES

1. T. Asano and D. Kirkpatrick, *Distance trisector curves in regular convex distance metrics*, Proceedings of the 3rd International Symposium on Voronoi Diagrams in Science and Engineering (ISVD 2006), pp. 8–17.
2. T. Asano, J. Matoušek, and T. Tokuyama, *The distance trisector curve*, Adv. Math. **212** (2007), 338–360, a preliminary version in STOC 2006, pp. 336-343.
3. ______, *Zone diagrams: Existence, uniqueness, and algorithmic challenge*, SIAM J. Comput. **37** (2007), 1182–1198, a preliminary version in SODA 2007, pp. 756-765.
4. T. Asano and T. Tokuyama, *Drawing equally-spaced curves between two points*, Abstracts of the 14th Annual Workshop on Computational Geometry with a Focus on Open Problems, 2004, pp. 24–25.
5. F. Aurenhammer, *Voronoi diagrams - a survey of a fundamental geometric data structure*, ACM Computing Surveys, vol. 3, 1991, pp. 345–405.

6. J. Chun, Y. Okada, and T. Tokuyama, *Distance trisector of segments and zone diagram of segments in a plane*, Proceedings of the 4th International Symposium on Voronoi Diagrams in Science and Engineering (ISVD 2007), pp. 66–73.
7. ______, *Distance trisector of a segment and a point*, Interdisciplinary Information Sciences **16** (2010), 119–125, a preliminary version in ISVD 2007, pp. 66–73.
8. S. C. de Biasi, B. Kalantari, and I. Kalantari, *Mollified zone diagrams and their computation*, Transactions on Computational Science **14** (2011), 31–59, a preliminary version in ISVD 2010, pp. 171–180.
9. J. Dugundji and A. Granas, *Fixed point theory*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2003.
10. K. Imai, A. Kawamura, J. Matoušek, D. Reem, and T. Tokuyama, *Distance k-sectors exist*, Computational Geometry: Theory and Applications **43** (2010), 713–720, preliminary versions in SoCG 2010, pp. 210-215, arXiv 0912.4164 (2009).
11. J. Jost, *Riemannian Geometry and Geometric Analysis*, Universitext, Springer-Verlag, Berlin, 1995.
12. A. Kawamura, J. Matoušek, and T. Tokuyama, *Zone diagrams in Euclidean spaces and in other normed spaces*, Mathematische Annalen, to appear. Preliminary versions in SoCG 2010, pp. 216-221, arXiv 0912.3016 (2009).
13. W. A. Kirk and B. Sims (eds.), *Handbook of metric fixed point theory*, Kluwer Academic Publishers, Dordrecht, 2001.
14. B. Knaster, *Un théorème sur les fonctions d'ensembles*, Ann. Soc. Polon. Math. **6** (1928), 133–134.
15. E. Kopecká, D. Reem, and S. Reich, *Zone diagrams in compact subsets of uniformly convex spaces*, Israel Journal of Mathematics **188** (2012), 1–23, preliminary versions in arXiv 1002.3583 (2010), CCCG 2010, pp. 17-20.
16. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces, II: Function spaces*, Springer, Berlin, 1979.
17. A. Okabe, B. Boots, K. Sugihara, and S. N. Chiu, *Spatial Tessellations: Concepts and Applications of Voronoi Diagrams*, second ed., Wiley Series in Probability and Statistics, John Wiley & Sons Ltd., Chichester, 2000, with a foreword by D. G. Kendall.
18. S. Prus, *Geometrical background of metric fixed point theory*, Handbook of Fixed Point Theory (W. A. Kirk and B. Sims, eds.), Kluwer Acad. Publ., Dordrecht, 2001, pp. 93–132.
19. D. Reem, *An algorithm for computing Voronoi diagrams of general generators in general normed spaces*, Proceedings of the sixth International Symposium on Voronoi Diagrams in Science and Engineering (ISVD 2009), pp. 144–152.
20. ______, *Voronoi and zone diagrams*, Ph.D. thesis, The Technion, Haifa, 2010.
21. ______, *The geometric stability of Voronoi diagrams with respect to small changes of the sites*, (2011), Complete version in arXiv 1103.4125, Extended abstract in SoCG 2011, pp. 254-263.
22. D. Reem and S. Reich, *Zone and double zone diagrams in abstract spaces*, Colloquium Mathematicum **115** (2009), 129–145, arXiv 0708.2668 (2007).
23. S. Reich and I. Shafrir, *Nonexpansive iterations in hyperbolic spaces*, Nonlinear Anal. **15** (1990), no. 6, 537–558.
24. M. Sharir and P. Agarwal, *Davenport-Schinzel sequences and their geometric applications*, Cambridge University Press, 1995.
25. A. Tarski, *A lattice-theoretical fixpoint theorem and its applications*, Pacific J. Math. **5** (1955), 285–309.

IMPA - Instituto Nacional de Matemática Pura e Aplicada, Estrada Dona Castorina 110, Jardim Botânico, CEP 22460-320, Rio de Janeiro, RJ, Brazil. Part of this work was done while the author was at the University of Haifa and the Technion - Israel Institute of Technology.

E-mail address: `dream@impa.br`